

Orthopaedic Surgery David Brown, MD Mary Joan Black, MD Michael W. Funderburk, MD Michael R. Henderson, MD Michael P. Hoenig, MD Chi Lim, MD

M. David Mitchell, MD Gerald L. Rollins, MD

Neurosurgery Christopher Chittum, MD Phillip Esce, MD Timothy Monroe,

Pain Management James Behr, MD

Anthony DiNicola, MD Jeffrey Smith, MD

Greater Trochanteric Bursitis

The greater trochanteric bursa is the fluid-filled sack that rests at the point where your muscle/tendons from your later lateral thigh and buttocks muscles attach to the hip bone. The bursa's function is to create a smooth surface across it, allowing the normal and painless movement. However, due to the repetitive movements of a certain body part, bursitis or inflammation of the bursa may occur. Symptoms of greater trochanteric bursitis include pain on the lateral side of your hip, difficulty sleeping on your side at night and pain while walking or climbing stairs. There are several effective treatment options for greater trochanteric bursitis including physical therapy, rest, anti-inflammatories, injections, and stretching exercises.

Trochanteric Bursitis Rehabilitation Exercises



North Grove Office

1330 Boiling Springs Road, Suite 1600

Court

Spartanburg, SC 29303 (864) 582-6396 Fax (864)542-2939 (864)721-0035

Neurosurgery Office 1075 Boiling Springs Road

Spartanburg, SC 29303 (864) 583-7265 Fax (864) 591-0422 West Grove 115 Deacon Tiller

Duncan, SC 29334 (864) 721-0025 Fax